

# **Dynamic Effects of Airborne Water Droplets on Air-Sea Interactions: Sea-Spray and Rain**

Fabrice Veron  
University of Delaware  
College of Marine Studies  
112C Robinson Hall  
Newark, DE 19716  
phone: (302) 831-6658 fax: (302) 831-6838 email: [fveron@udel.edu](mailto:fveron@udel.edu)

Award Number: N00014-05-1-0609

## **LONG-TERM GOALS**

We anticipate that this research will lead to an improved understanding of water droplet mediated air-sea fluxes with implications in basic research, satellite or land-based remote sensing of the ocean surface, electromagnetic communication, military surveillance, and weather forecasting. We foresee that this work will become the initial stages of a long-term effort involving further development of theoretical work, and laboratory and field experiments.

## **OBJECTIVES**

In recent years, small-scale surface dynamics (waves, turbulence, bubbles, and drops) have been recognized as crucial to global climate through their impact on air-sea fluxes. However, the dynamics of a turbulent surface can be quite complicated, and at times (at high wind speed) the main problem is to identify the surface which then resembles a mixture of bubbles and drops. While the upper ocean bubbly mixture has been relatively well studied, especially in the context of the acoustics, it appears that airborne water droplets have been widely ignored despite their undeniable importance.

Recent work point toward the fact that airborne water droplets can exchange with the atmosphere a consequent amount of momentum, energy, and enthalpy, especially at high wind speeds (Andreas et al., 1995; Andreas, 2004). At low wind speed, it appears that a significant fraction of shear stress can be carried by falling rain and transmitted to the wind field (Caldwell and Elliot, 1971). The direction and amplitude of the momentum and enthalpy fluxes depend on the origin of the droplet (the upper atmosphere for the rain, the ocean surface for sea spray), and in the case of sea spray, if it evaporates entirely or falls back.

This project is aimed at improving and use an existing sea-spray Lagrangian turbulent transport and evaporation model to study the heat, moisture and momentum balances in the lower atmospheric boundary layer when water droplets are present. We also plan to conduct laboratory experiments to assess the effects of impacting rain on the surface and the consequent exchange of energy from the wind to the water via the droplets.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>SEP 2007</b>		2. REPORT TYPE <b>Annual</b>		3. DATES COVERED <b>00-00-2007 to 00-00-2007</b>	
4. TITLE AND SUBTITLE <b>Dynamic Effects Of Airborne Water Droplets On Air-Sea Interactions: Sea-Spray And Rain</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>University of Delaware, College of Marine Studies, 112C Robinson Hall, Newark, DE, 19716</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>code 1 only</b>					
14. ABSTRACT <b>We anticipate that this research will lead to an improved understanding of water droplet mediated air-sea fluxes with implications in basic research, satellite or land-based remote sensing of the ocean surface, electromagnetic communication, military surveillance, and weather forecasting. We foresee that this work will become the initial stages of a long-term effort involving further development of theoretical work, and laboratory and field experiments.</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>6</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## **APPROACH**


We propose to improve our current sea-spray Lagrangian turbulent transport and evaporation model to include several important effects that have been neglected in the past. Also, we propose to generalize part of the parameterizations such that the model is adequate for a wide range of conditions and therefore usable to study both suspended sea-spray and also rain droplets. The numerical part of this project will be aimed at further developing the model, and making use of a cluster of PC computers to run the calculations. In addition, a laboratory experimental part is proposed for the later years, in which we will examine the momentum exchanges between impacting rain and the ocean surface.

## **WORK COMPLETED**

We have implemented a large computer system consisting of twenty-four, single processor PCs on which the model processing is performed. The data are then recombined on a large server computer for analysis. We have modified the currently existing model to account for anisotropy in the Lagrangian turbulence in the air-flow above the wavy ocean surface, and include wave-turbulence interactions. This has been done via a generalization of the Lagrangian Stochastic equations for turbulent fields (Wilson and Sawford 1996). The generalization includes non-isotropic conditions allowing us to incorporate stratification effects in both the mean Eulerian, and non-homogenous conditions allowing difference in statistical properties of the turbulence upwind and downwind of surface waves. This is critical in particular for sea-spray suspension dynamics where the spray is generated on the front (leeward) face of the wave. To date, we believe we have successfully implemented turbulence closures and the wave effects. Turbulent for the passive scalars has also been included. Furthermore, we have developed a general description of the wind, temperature and humidity velocity profiles above the wavy surface which includes the diffusive molecular layer based on hybrid equations from the classical Van Driest (viscous) and logarithmic layers. The full wave spectrum also generates a moving bottom boundary which then interacts with the mean profiles as well as the turbulence. Based on the wave spectrum and breaking statistics, we have also incorporated the effects of air-flow separation on the momentum flux. These effects feed into the determination of the viscous and turbulent partition and thereby influence the calculation of the wind profiles. This leads to a non-linear formulation for the sea surface stress. To date, we have completed coding of the Lagrangian model and have performed preliminary runs with full droplets size distribution that will serve as debugging trials and will generate preliminary data. In addition, we have initiated the experiments on the effect of impacting rain on the surface and the consequent transport of kinetic energy from the wind to the surface. Our preliminary results show that significant kinetic energy is injected in the surface layers of the water and that penetration depth and rate are functions of the rain rate.

## **RESULTS**


We have developed a Spray Lagrangian Turbulent Transport and Evaporation model (hereafter SpLaTTE) which includes droplet microphysics, propagating surface waves, atmospheric stratification, turbulence, and the complete linear unsteady equation of motion for the droplets. Our results indicate that the short thermal and inertial response time of small droplets is such that they decelerate and warm up significantly as they approach the thermal and momentum boundary layer near the ocean surface. We have therefore spent considerable effort in developing a generalized form for the wind and scalar profiles (including stability), which account for the viscous layer close to the interface. Figure 1 shows



**Figure 1. Wind profiles from 2m/s to 40m/s in 2m/s increment and plotted in wall layer coordinate. Note the presence of a well defined viscous sub-layer and the effect of stratification at low wind speed (the top most profiles for large  $z$ ).**

the wind profile above the surface (plotted in wall layer coordinate) for wind speeds from 2 to 40m/s in increments of 2m/s. The departure from the log-layer at the lowest wind speed is due to effects of atmospheric stratification, effects that disappear at higher wind speeds. Another significant improvement to the model was the inclusion of fully non-isotropic Lagrangian turbulence for the velocity, temperature and humidity variables. Figure 2 shows spectra for the vertical velocity of both a 150 $\mu$ m droplet and the air velocity around the droplet. In order to assure a sufficiently long residence time to obtain statistical properties for this test case, the droplet was released at 10 meters above the wavy bottom boundary. We found that the air velocity around the droplet is not significantly different from the true Lagrangian velocity that would be experienced by a fluid particle. Because the air velocity at the droplet is decorrelated in both time and separation from an idealized fluid trajectory, the spectrum is slightly flatter than the theoretical  $\omega^{-2}$  behavior for a fluid Lagrangian spectrum. The spectrum of the drop velocity, however, shows an  $\omega^{-4}$  dependence as is expected. Figure 2 shows that at lower frequencies, the particle and air at the particle are in phase and then move out of phase around the stokes response frequency, 14.3 Hz in this case. The results in figure 2 suggest that our model correctly simulates the trajectory of a particle in a Lagrangian turbulent airflow. It should be noted here that the simulated turbulence will uniformly have an approximate  $\omega^{-2}$  dependence, and therefore, the very high frequencies at which a dissipation range should exist will not be accurately modeled. But considering that the spectrum for the particle velocity follows a  $\omega^{-4}$  law, the contribution of very high frequencies to the total particle velocity variance is expected to be negligible. In addition, it should also be noted that the particle moves through the atmospheric surface boundary layer and that consequently, the nature and strength of the turbulence will also change as the particle, for example, changes altitude. As a result, since individual droplets are followed through their course in the boundary layer, we did not expect to obtain an  $\omega^{-2}$  dependence in the air-velocity spectrum surrounding a droplet along its path. Finally, as the particle stays suspended in the air-flow, its density

and radius will change which will affect its response time and behavior vis-à-vis the forcing from the turbulence in the air during the




**Figure 2. (a) Lagrangian frequency spectra for the vertical air velocity at the particle location (black), and for the vertical droplet velocity (gray), (b) phase relationship between the air and particle velocities, and (c) the coherence squared values between the air and particle velocities.**

entirety of its flight. This will in turn render spectral analysis along the flight rather difficult to interpret. We feel however, that this is one of the strength of the Lagrangian approach since we can keep track of both the droplet and the atmospheric properties surrounding it at all times.


We believe we have successfully implemented the closures and wave effects in the simulation of the Lagrangian turbulence. Finally, we have also incorporated the effects of air-flow separation on the full air-sea momentum balance. The air-flow separation is based on breaking wave statistics (in turn based on the balance of wave growth and dissipation), and on the physics of the detachment over back facing steps. The air-flow separation past a breaking wave reduces the viscous stress in the recirculating bubble and also enhances turbulence thereby affecting the partition of air-sea stress into its viscous, turbulent, and wave-induced components. This new parameterization of the air-sea stress amounts to a non-linear formulation as the air-flow separation feeds on other stress components, but also on the determination of the mean velocity profile, which in turn affect the wave spectrum. This sub-model reproduces the observed features of the drag coefficient from low to high wind speeds despite extrapolating empirical wave spectra and breaking wave statistics beyond known limits. The model shows the saturation of the drag coefficient at high wind speeds for both field and laboratory fetches, suggesting that air-flow separation over ocean waves and its accompanying effects may play a significant role in the driving physics of the air-sea stress, at least at high wind speeds (figure 3).

In the laboratory, we have initiated the study of the impact on the rain on the air-sea momentum flux. Preliminary tests have been conducted in which we found that the rain can significantly alters the wind speed profiles as momentum is exchanged between the two phases (air and airborne water). As the rain droplets fall they decelerate while approaching the surface, thereby accelerating the wind in the near-surface layers. Impact velocity is approximately 85% of the 10-m wind speed. It should be noted that this effect also exist with sea spray, albeit to a lesser extent because of the size/mass difference. Also, the momentum exchange is in the opposite direction as spray significantly accelerates while airborne, thereby extracting momentum from the airflow. Upon impact, the rain transmits its kinetic energy to the water in the form of turbulence and currents. Figure 4 below shows turbulent kinetic energy

profiles up to depths of approximately 15cm (under no wind conditions) as the rain starts falling of the water column. Our results indicate that the penetration depth, as well as the peneration speed is dependent on the rain rate.



**Figure 3. a) Drag coefficient as a function of wind speed for 10 km (solid), 100 km fetch (dash-dotted), and infinite (dotted) fetches along with the experimental data of Large and Pond (1981) (light gray), Taylor and Yelland (2000) (dark gray) and Powell et al. (2003) (gray symbols). Model results from Kudryavtsev and Makin (2007) (dashed line) are also included. b) Fraction of the viscous (line), wave-induced (dashed), and separation (dash-dotted) stresses as a function of wind speed for a 100 km fetch**



**Figure 4. Profile of the turbulent Kinetic Energy under the inception of the rain, for different rain rate conditions**

## IMPACT/APPLICATIONS

We anticipate that this research will lead to an improved understanding of water droplet mediated air-sea fluxes. Potential impacts are cited above and also include a better understanding of air-sea fluxes, especially at high wind speeds when significant amount of water droplets can be suspended in the air-flow. In particular in storm conditions or hurricanes, there can be a combined effect of sea-spray and rain which then leads to competing droplet-mediated momentum fluxes. The spray is expected to carry some momentum from the ocean to the atmosphere, while the rain will impart momentum to the ocean surface. We anticipate that our model will be allowing us to determine the net momentum fluxes carried by different droplet types. In addition, the model should lead to heat and moisture fluxes as well, which are a critical component of storm dynamics.

## REFERENCES

Andreas, E. L., J. B. Edson, E. C. Monahan, M. P. Rouault, and S. D. Smith, 1995: The spray contribution to net evaporation from the sea: a review of recent progress. *Bound.-Layer Meteor.*, 72,3-52.

Andreas, E. L., 2004: Spray stress revisited. *J. Phys. Oceanogr.*, 34,1429-1440.

Caldwell, D. R., and W. P. Elliott, 1971: Surface stress produced by rainfall. *J. Phys. Oceanogr.*, 1,145-148.

Kudryavtsev, V.N. and V.K. Makin, 2007: Aerodynamic roughness of the sea surface at high winds. *Bound.-Layer Meteorol.*, doi:10.1007/s10546-007-9184-7

Large, W.G. and S. Pond, 1981: Open ocean momentum flux measurements in moderate to strong winds, *J. Phys. Oceanogr.*, 11,324-336

Taylor, P.K. and M.J. Yelland, 2000: On the apparent “‘imbalance” term in the turbulent kinetic energy budget. *J. Atmos. Ocean. Tech.*, 17,82-89

Powell, M.D., P.J. Vickery, and T.A. Reinhold, 2003: Reduced drag coefficient for high wind speeds in tropical cyclones. *Nature*, 422, 279-283

Wilson, J.D. and B.L. Sawford, 1996: Review of Lagrangian Stochastic models for trajectories in the turbulent atmosphere. *Bound. Layer Meteor.*, 78, 191-210.

## PUBLICATIONS

Mueller J. and F. Veron, 2007: Nonlinear formulation of the bulk surface stress over breaking waves: a simple feedback mechanism resulting from air-flow separation, *Bound.- Layer Meteor.*

Mueller J. and F. Veron, 2007: Dispersive transport model for heavy particles within the marine boundary layer, *Bound.- Layer Meteor*